

Stockholm
2007

NOTE ON NON-METRIC GRAVITY

Ingemar Bengtsson¹

*Stockholm University, AlbaNova
Fysikum
S-106 91 Stockholm, Sweden*

Abstract

We discuss a class of alternative gravity theories that are specific to four dimensions, do not introduce new degrees of freedom, and come with a physical motivation. In particular we sketch their Hamiltonian formulation, and their relation with some earlier constructions.

¹Email address: ingemar@physto.se. Supported by VR.

A class of alternative gravity theories were recently introduced, under the name of “non-metric gravity” [1]. There are reasons to take this class seriously, in particular arguments were advanced why this class—which is defined by one free function of two variables—should be closed under renormalization [2]. Indeed the construction is interesting already on the classical level, since it does not introduce any new degrees of freedom, as compared to GR (which is a member of the class); moreover the construction is intrinsically four dimensional (arguably a good thing), and is based on a clean split between conformal structure and conformal factor of the metric. In fact the split is so clean that the latter is largely lost track of, which brings us to some weak points of these models. The first of those is that the models describe complex spacetimes, and it is not clear how to recover the Lorentzian sector. The second is that they appear to be quite difficult to couple to matter (but this may turn out to be a strong point in the end).

The purpose of this note is to clarify the relation between non-metric gravity and an earlier class of models with similar properties [3, 4]. The starting point is Plebański’s action for vacuum general relativity (GR) [5, 6, 7]:

$$S = \frac{1}{4} \int d^4x \, \epsilon^{\alpha\beta\gamma\delta} \left[\Sigma_{\alpha\beta i} F_{\gamma\delta i} - \frac{1}{2} \Phi_{ij} \Sigma_{\alpha\beta i} \Sigma_{\gamma\delta j} \right] , \quad (1)$$

where Φ is a symmetric traceless three by three matrix,

$$F_{\alpha\beta i} = \partial_\alpha A_{\beta i} - \partial_\beta A_{\alpha i} + i f_{ijk} A_{\alpha j} A_{\beta k} , \quad (2)$$

and f_{ijk} are the structure constants of the $SU(2)$ Lie algebra. To discuss the corresponding space of solutions, introduce a spacetime metric given in terms of a tetrad of vector fields by

$$g_{\alpha\beta} = \eta_{IJ} e_\alpha^I e_\beta^J , \quad (3)$$

where η_{IJ} is the Minkowski metric. Use the tetrad to form a 2-form

$$\sigma_{\alpha\beta}^{IJ} = e_{[\alpha}^I e_{\beta]}^J . \quad (4)$$

One can show that, in any solution of the equations coming from Plebański’s action, the 2-form $\Sigma_{\alpha\beta i}$ is the self dual part of a 2-form $\sigma_{\alpha\beta}^{IJ}$, the metric $g_{\alpha\beta}$ solves Einstein’s vacuum equations, the connection $A_{\alpha i}$ is the metric compatible self dual spin connection, and Φ_{ij} is the self dual part of the

Weyl tensor—which is indeed naturally a symmetric traceless complex three by three matrix [8], even if it is more common to regard it as a rank 4 spinor. We will not explain these matters further here—the argument given below will lead to the same conclusion in a different way.

Krasnov modifies this action by setting

$$\Phi_{ij} = \phi_{ij} + \delta_{ij}\Lambda(\text{Tr}\phi^2, \text{Tr}\phi^3) , \quad (5)$$

where $\text{Tr}\phi = 0$. If the function Λ is constant this results in Einstein's theory including a cosmological constant. Otherwise we have a new class of models, not equivalent to GR.

Although we are motivated by our belief that Krasnov's construction is connected to some deep mathematical properties of four dimensional manifolds, the only mathematics we need for now is the characteristic equation for three by three matrices, namely

$$M^3 - M^2\text{Tr}M - \frac{1}{2}M(\text{Tr}M^2 - (\text{Tr}M)^2) - \det M = 0 . \quad (6)$$

It follows that

$$\text{Tr}M^3 - \frac{3}{2}\text{Tr}M^2\text{Tr}M + \frac{1}{2}(\text{Tr}M)^3 - 3\det M = 0 . \quad (7)$$

A three by three matrix has three independent invariants only. Thus equipped, we can proceed to cast Krasnov's models into Hamiltonian form. Afterwards we will make some further comments, in particular we will discuss how a spacetime metric enters the game.

We begin. Vary the action (1) with respect to the 2-form $\Sigma_{\alpha\beta i}$, solve for the same, and insert the result back into the action. The result is

$$S = \frac{1}{8} \int d^4x \epsilon^{\alpha\beta\gamma\delta} \Psi_{ij} F_{\alpha\beta i} F_{\gamma\delta j} , \quad (8)$$

where

$$\Psi \equiv \Phi^{-1} = \frac{3(\phi^2 - \phi\Lambda - \frac{1}{2}\text{Tr}\phi^2 + \Lambda^2)}{\text{Tr}\phi^3 - \frac{3}{2}\text{Tr}\phi^2\Lambda + 3\Lambda^3} . \quad (9)$$

Here we used the characteristic equation, and we closed our eyes to the fact that Φ may be non-invertible—this happens when its algebraic Petrov type is $\{-\}$, $\{4\}$, or $\{3, 1\}$. The invariants of the matrix Ψ are given by

$$I_1 \equiv (\text{Tr}\Psi)^2 - \text{Tr}\Psi^2 = \frac{18\Lambda}{\text{Tr}\phi^3 - \frac{3}{2}\text{Tr}\phi^2\Lambda + 3\Lambda^3} \quad (10)$$

$$I_2 \equiv \text{Tr}\Psi = \frac{3(3\Lambda^2 - \frac{1}{2}\text{Tr}\phi^2)}{\text{Tr}\phi^3 - \frac{3}{2}\text{Tr}\phi^2\Lambda + 3\Lambda^3} \quad (11)$$

$$I_3 \equiv \det\Psi = \frac{3}{\text{Tr}\phi^3 - \frac{3}{2}\text{Tr}\phi^2\Lambda + 3\Lambda^3} . \quad (12)$$

Since $\Lambda = \Lambda(\text{Tr}\phi^2, \text{Tr}\phi^3)$, these are three functions of two variables. Hence there will exist a constraint of the form

$$h(I_1, I_2, I_3) = 0 . \quad (13)$$

The form of the function h will depend on the form of the function Λ . For the moment we pass over some technical difficulties that arise at this point.

Coming back to the action (8), we perform a 3+1 split through $\alpha = (t, a)$. We define the “magnetic” field

$$B_i^a \equiv \frac{1}{2}\epsilon^{tabc}F_{bci} \equiv \frac{1}{2}\epsilon^{abc}F_{bci} . \quad (14)$$

We denote time derivatives with overdots. Then the canonical momentum is

$$E_i^a \equiv \frac{\partial\mathcal{L}}{\partial\dot{A}_{ai}} = \Psi_{ij}B_j^a . \quad (15)$$

One finds that the canonical Hamiltonian vanishes. Variation with respect to the traceless matrix ϕ , in terms of which the matrix Ψ is defined, gives a consistency condition which is automatically obeyed, given the constraints. Primary constraints arise because the matrix Ψ is symmetric and constrained by eq. (13). The first primary constraint is

$$\Psi_{ij} = \Psi_{ji} \quad \Leftrightarrow \quad \mathcal{H}_a \equiv \epsilon_{abc}E_j^b B_k^c \approx 0 . \quad (16)$$

To formulate the second primary constraint, let us introduce some further notation. Define

$$EEB \equiv \epsilon_{abc}f_{ijk}E_i^a E_j^b B_k^c = ((\text{Tr}\Psi)^2 - \text{Tr}\Psi^2) \det B \equiv \frac{1}{6}I_1 BBB . \quad (17)$$

In a similar notation

$$EBB = \frac{1}{3}I_2 BBB \quad EEE = I_3 BBB . \quad (18)$$

Then one finds

$$h(I_1, I_2, I_3) = 0 \quad \Leftrightarrow \quad \mathcal{H}(EEE, EEB, EBB, BBB) \approx 0 , \quad (19)$$

where $\mathcal{H}$ is a homogeneous function of order 1. Finally a secondary constraint will arise when we vary the action with respect to A_{ti} . The 3+1 decomposition is now complete, and we have arrived at the phase space action

$$S = \int d^4x \dot{A}_{ti} E_i^a - N\mathcal{H} - N^a \mathcal{H}_a + A_{ti} D_a E_i^a . \quad (20)$$

Lagrange multipliers were introduced for the primary constraints. We have chosen a density weight equal to -1 for the lapse function N .

Now vary the phase space action with respect to E_i^a . In most cases one can solve the resulting equation for the same variable, and insert the solution back into the phase space action. With the definitions

$$\eta \equiv \frac{3i}{8N BBB} \quad (21)$$

$$\Omega_{ij} \equiv \epsilon^{\alpha\beta\gamma\delta} F_{\alpha\beta i} F_{\gamma\delta j} \quad (22)$$

(yet another three by three matrix!), one arrives in this way at the CDJ type [9, 3] action

$$S = \frac{1}{8} \int d^4x L(\text{Tr}\Omega, \text{Tr}\Omega^2, \text{Tr}\Omega^3, \eta^{-1}) , \quad (23)$$

where again the function L is homogeneous of order 1. In the calculation one must solve for the invariants of Ψ as functions of the invariants of Ω ; in practice this may be difficult. If we impose the constraint on Ψ through a Lagrange multiplier η , the CDJ action can be arrived at directly from the action (8).

This ends our overview of the equivalence between the Plebański, CDJ, and Hamiltonian formulations of the models considered by Krasnov. We have hidden some difficulties which will appear when we consider a few examples. Let us start from the action (1). In eq. (5) we set

$$\Lambda = i\lambda/3 , \quad (24)$$

where λ is a constant. It follows that $I_1 = 2i\lambda I_3$, and the Hamiltonian constraint is

$$\mathcal{H} = \frac{i}{2}EEB + \frac{\lambda}{3}EEE . \quad (25)$$

This is Ashtekar's Hamiltonian for GR including a cosmological constant [10] (and we included a constant factor for agreement with standard conventions). This establishes the equivalence between the original Plebański action and the Einstein-Hilbert action. The corresponding CDJ action is quite elegant when $\lambda = 0$ (see below), but otherwise one encounters a difficulty. To solve for the invariants of Ψ in terms of the invariants of Ω one must take a square root. Consequently the CDJ action will have two branches—not to mention the fact that it becomes very complicated [11].

In our second example we let the matrix Ψ have constant trace. Starting from the action (8) it is easily deduced that

$$\text{Tr}\Psi = -i\alpha \quad \Rightarrow \quad \mathcal{H} = \frac{1}{2}EBB + \frac{i\alpha}{6}BBB . \quad (26)$$

This theory is “dual” to Ashtekar's. The parameter α can be set to zero through the canonical transformation

$$A_{ai} \rightarrow A_{ai} \quad E_i^a \rightarrow E_i^a - \frac{i\alpha}{3}B_i^a . \quad (27)$$

Let us do so for simplicity. It is clear from eq. (11) that there are two possibilities for deriving this model from Krasnov's action, namely

$$\Lambda = \pm \sqrt{\text{Tr}\phi^2/6} . \quad (28)$$

The CDJ action does not exist in this case, since the equation obtained by varying the phase space action with respect to E_i^a is independent of E_i^a .

Our third and final example is a one-parameter family of models that attracted some attention because of its simple CDJ Lagrangian [12, 13]:

$$S = \frac{1}{8} \int d^4x \, \eta \left(\text{Tr}\Omega^2 + a(\text{Tr}\Omega)^2 \right) . \quad (29)$$

The particular case $a = -1/2$ gives standard GR with vanishing cosmological constant. The Hamiltonian constraint becomes

$$\mathcal{H} = \frac{i}{2} EEB - \frac{3i(1+2a)}{4(1+3a)} \frac{(EBB)^2}{BBB} . \quad (30)$$

Hence

$$\mathcal{H} = 0 \quad \Leftrightarrow \quad I_1 - \frac{1+2a}{1+3a} I_2 = 0 \quad (31)$$

Using eqs. (10-11) we find a quadratic equation for the function Λ , namely

$$6\Lambda = \frac{1+2a}{1+3a} \left(3\Lambda^2 - \frac{1}{2} \text{Tr}\phi^2 \right) . \quad (32)$$

Consequently Krasnov's action will again have two branches.

We have now explained the practical difficulties involved in carrying out the steps relating the Krasnov, CDJ, and phase space actions. But where is the spacetime metric in all this? Krasnov's answer is that the algebraic constraint on $\Sigma_{\alpha\beta i}$, obtained by varying the action (5) with respect to ϕ , will enable one to introduce a tetrad and a metric also in the case when $\Phi(\phi)$ is not traceless. However, the resulting metric is defined only up to an arbitrary conformal factor, hence the name “non-metric gravity”.

The Hamiltonian formulation opens another avenue to the same end. The constraint algebra will include the bracket

$$\{\mathcal{H}[N], \mathcal{H}[M]\} = \mathcal{H}_a[(N\partial_a M - M\partial_a N)q^{ab}] , \quad (33)$$

where $\mathcal{H}[N]$ means that the constraint has been smeared with a test function of the appropriate density weight. The tensor density q^{ab} will be a definite function of the canonical variables E_i^a and B_i^a . In order to agree with the usual geometric interpretation of the constraint algebra [14], we must set

$$gg^{ab} \equiv q^{ab} , \quad (34)$$

where g_{ab} is the spatial metric. If one then runs the ADM decomposition of the spacetime metric backwards, one finds (after a non-trivial calculation [15]) that the latter must be

$$g_{\alpha\beta} = \frac{4}{3} \eta f_{ijk} \epsilon^{\mu\nu\rho\sigma} F_{\alpha\mu i} F_{\nu\rho j} F_{\sigma\beta k} . \quad (35)$$

This is an interesting expression, since it actually implies that the 2-forms $F_{\alpha\beta i}$ are self dual with respect to this metric—and conversely [16, 17]. Since Krasnov comes to the same conclusion, the two ways of defining the conformal structure agree. It should be noted that the conformal factor adopted in eq. (35) is conventional also from the Hamiltonian perspective, because one can redefine the lapse function with scalar functions of the canonical variables without changing the dynamical content of the theory—this would change the tensor density q^{ab} in eq. (33) with a factor, and hence the conformal factor of the metric.

If these models are to be taken seriously as alternative gravity theories, it must be properly explained how to make the restriction to real Lorentzian spacetimes, and it must be shown how to couple them to matter degrees of freedom. Krasnov’s formulation is likely to be helpful here. If it can breathe new life into the asymptotic safety programme [18], it is so much the better.

Acknowledgement

I thank Kirill Krasnov for several interesting emails.

References

- [1] K. Krasnov, *Non-Metric Gravity I: Field Equations*, arXiv eprint gr-qc/0307002.
- [2] K. Krasnov, *Renormalizable Non-Metric Quantum Gravity?*, arXiv eprint hep-th/0611182.
- [3] P. Peldán, *Actions for gravity, with generalizations: A review*, Class. Quant. Grav. **11** (1994) 1087.
- [4] I. Bengtsson, *Clifford algebra of two-forms, conformal structures, and field equations*, in Z. Oziewicz et al. (eds.): *Spinors, Twistors, Clifford Algebras and Quantum Deformations*, Kluwer 1993. Also available as gr-qc/9210001.
- [5] J. F. Plebański, *On the separation of Einsteinian substructures*, J. Math. Phys. **18** (1977) 2511.

- [6] R. Capovilla, J. Dell, T. Jacobson and L. Mason, *Self-dual 2-forms and gravity*, Class. Quant. Grav. **8** (1991) 41.
- [7] M. Reisenberger, *New constraints for canonical general relativity*, Nucl. Phys. **B457** (1995) 643.
- [8] J. Ehlers and W. Kundt, *Exact solutions of the gravitational field equations*, in L. Witten (ed.): Gravitation: an introduction to current research, Wiley 1962.
- [9] R. Capovilla, J. Dell and T. Jacobson, *A pure spin connection formulation of gravity*, Class. Quant. Grav. **8** (1991) 59, **9** (1992) 1839 (E).
- [10] A. Ashtekar, *New Hamiltonian formulation of general relativity*, Phys. Rev. **D36** (1987) 1587.
- [11] P. Peldán, *Legendre transforms in Ashtekar's theory of gravity*, Class. Quant. Grav. **8** (1991) 1765.
- [12] R. Capovilla, *Generally covariant gauge theories*, Nucl. Phys. **B373** (1992) 233.
- [13] I. Bengtsson and P. Peldán, *Another "cosmological" constant*, Int. J. Mod. Phys. **A7** (1992) 1287.
- [14] S. Hojman, K. Kuchař and C. Teitelboim, *Geometrodynamics regained*, Ann. of Phys. **96** (1976) 88.
- [15] I. Bengtsson, *Self-duality and the metric in a family of neighbours of Einstein's equations*, J. Math. Phys. **32** (1991) 3158.
- [16] M. Schönberg, *Electromagnetism and gravitation*, Rev. Bras. Fis. **1** (1971) 91.
- [17] H. Urbantke, *On integrability properties of $SU(2)$ Yang-Mills fields. I. Infinitesimal part*, J. Math. Phys. **25** (1984) 2321.
- [18] S. Weinberg, *Ultraviolet divergences in quantum theories of gravitation*, in S. W. Hawking and W. Israel: General Relativity. An Einstein Centenary Survey, Cambridge UP 1979.